

*OB-GYN SPECIALISTS
WOMEN CARING FOR WOMEN*

2121 Ygnacio Valley Rd., E101
Walnut Creek, CA 94598
Tel: (925) 945-6600
Fax: (925) 945-7842

Judith Hartman, MD
Janine Senior, MD
Janette Walker, MD
Sari Kasper, DO
Heather West, MS., ANP
Linda C. Wight, RNC, WHCNP

Post-LEEP Instructions

What do I need to know as I recover at home?

You may feel tired or weak, and it is best if you do not go home alone

You may have cramps for a short time which should be lessened with ibuprofen (Advil) or Alleve.

Please do not swim or take a tub bath for 48 hours. Showering is okay.

You may have vaginal discharge for 2-3 weeks, ranging in color from pink to black. You may also have bleeding similar to a period.

You should not have intercourse or use tampons until the cervix heals. This takes about 2-3 weeks.

If you have vaginal discharge or bleeding after 2-3 weeks, give yourself longer to heal and refrain from putting anything into the vagina.

Avoid heavy lifting or vigorous exercise for 2-3 weeks.

Call our office immediately for:

Heavy or increased vaginal bleeding that soaks more than one pad per hour for 1-2 hours

Dizziness or fainting

Chills or fever of greater than 101

Follow up Care

Please schedule a two week follow-up visit to evaluate healing

The abnormal cells on your cervix may return and it is important to follow your doctor's instructions for repeat pap smears in the future.